

National Air Quality and Emissions Trends Report, 1996

U.S. Environmental Protection Agency
Office of Air Quality Planning and Standards
Emissions Monitoring and Analysis Division
Air Quality Trends Analysis Group
Research Triangle Park, North Carolina 27711

January 1998

About the Cover

The cover provides a visual air quality comparison of the average best and worst visibility days at Great Smoky Mountain National Park from 1992 to 1995. The image was generated using software called WinHaze. WinHaze, developed by Air Resource Specialists of Fort Collins, Colorado, uses visual range parameters to degrade a pristine image, thus simulating what a scene would look like with the given visibility parameters. Images such as these are helpful in defining and communicating the visibility problem and assessing any progress made. Additional information on visibility can be found in Chapter 3 of this report.

Disclaimer

This report has been reviewed and approved for publication by the U.S. Environmental Protection Agency's Office of Air Quality Planning and Standards. Mention of trade names or commercial products are not intended to constitute endorsement or recommendation for use.

Acknowledgments

The Trends Team would like to acknowledge Kate Ramoth of GeoLogics Corporation for assistance with layout, tables, graphics, and technical editing; the parties who reviewed this report prior to publication for their comments; and the following individuals for their extensive contributions in a variety of areas: Dr. John Ackermann, John Bachmann, Angela Bandemehr, Desmond Bailey, Dr. Jane Caldwell, Rich Cook, William Cox, Rich Damberg, Barbara Driscoll, Kathy Kaufmann, Mary Manners, Dr. Karen Martin, Melissa McCullough, Dr. Dave McKee, David Misenheimer, Dr. Diedre Murphy, Sharon Nizich, Anne Pope, Kelly Rimer, Dr. Mary Ross, Dr. Roy Smith, Greg Stella, Lori Stewart, and Dr. Al Wehe.

Preface

This is the twenty-fourth annual report on air pollution trends in the United States issued by the U.S. Environmental Protection Agency. The report is prepared by the Air Quality Trends Analysis Group (AQTAG) in Research Triangle Park, North Carolina and is directed toward both the technical air pollution audience and other interested parties and individuals.

The report, complete with graphics and data tables, can be accessed via the Internet at <http://www.epa.gov/oar/aqtrnd96/>. AQTAG solicits comments on this report and welcomes suggestions regarding techniques, interpretations, conclusions, or methods of presentation. Comments can be submitted via the website or mailed to:

Attn: Trends Team
AQTAG (MD-14)
U.S. EPA
Research Triangle Park, NC 27711

For additional air quality data, readers can access the Aerometric Information Retrieval System's (AIRS) executive software at <http://www.epa.gov/oar/airs/aewin>.

Contents

CHAPTER 1	
Executive Summary	1
Overview and Highlights	1
Improvement in the Face of Economic Growth	2
The Need for Continued Progress	3
References	4
CHAPTER 2	
Air Quality Trends	7
Carbon Monoxide	9
Lead	13
Nitrogen Dioxide	17
Ozone	21
The New 8-hour Ozone Standards	27
Determining Compliance with the New 8-hour Ozone Standards	28
Particulate Matter	30
The New Particulate Matter Standards	34
Determining Compliance With the New PM Standards	35
Sulfur Dioxide	38
References	42
CHAPTER 3	
Visibility Trends	43
Introduction	43
Nature and Sources of the Problem	43
Long-Term Trends	45
Recent Trends in Rural Areas: 1988-1995	45
Regional Trends	46
Current Conditions	48
Programs to Improve Visibility	49
References	52
CHAPTER 4	
PAMS: Enhanced Ozone & Precursor Monitoring	53
Background	53
Network Requirements	53
Monitoring Requirements	54
Program Objectives	55
VOC Characterization	55

CHAPTER 4	
PAMS: Enhanced Ozone & Precursor Monitoring (continued)	
Trends	55
NO _x Versus VOC	57
Summary	58
References	58
CHAPTER 5	
Air Toxics.....	61
Background.....	61
Ambient Air Quality Data	64
Air Toxics Control Program	65
Air Toxics Regulation and Implementation Status	66
Emissions Reductions Through the MACT Program	67
Residual Risk	67
Special Studies/Programs	67
References	70
CHAPTER 6	
Nonattainment Areas	73
CHAPTER 7	
Metropolitan Area Trends	75
Status: 1996	75
Trends Analysis	75
The Pollutant Standards Index	76
Summary of PSI Analyses	76
References	78
APPENDIX A	
Data Tables	79
APPENDIX B	
Methodology	149
Air Quality Data Base	149
Air Quality Trend Statistics	150
References	152

Figures

Figure 1-1. Total U.S. population, vehicle miles traveled, U.S. gross domestic product, and aggregate emissions, 1970–1996.	3
Figure 1-2. Number of people living in counties with air quality concentrations above the level of the NAAQS in 1996.	3
Figure 2-1. Trend in second maximum non-overlapping 8-hour average CO concentrations, 1987–1996.	9
Figure 2-2. National total CO emissions trend, 1987–1996.	10
Figure 2-3. CO emissions by source category, 1996.	10
Figure 2-4. CO second maximum 8-hour concentration trends by location, 1987–1996.	11
Figure 2-5. Highest CO second maximum 8-hour concentration by county, 1996.	11
Figure 2-6. Long-term ambient CO trend, 1977–1996.	12
Figure 2-7. Trend in maximum quarterly average Pb concentrations (excluding source-oriented sites), 1987–1996.	13
Figure 2-8. National total Pb emissions trend, 1987–1996.	14
Figure 2-9. Pb maximum quarterly mean concentration trends by location (excluding source-oriented sites), 1987–1996.	14
Figure 2-10. Long-term ambient Pb trend, 1977–1996.	15
Figure 2-11. Pb emissions by source category, 1996.	15
Figure 2-12. Pb maximum quarterly concentration in the vicinity of Pb point sources, 1996.	16
Figure 2-13. Highest Pb maximum quarterly mean by county, 1996.	16
Figure 2-14. Trend in annual NO ₂ concentrations, 1987–1996.	17
Figure 2-15. National total NO _x emissions trend, 1987–1996.	18
Figure 2-16. NO _x emissions by source category, 1996.	18
Figure 2-17. NO ₂ annual mean concentration trend by location, 1987–1996.	19
Figure 2-18. Highest NO ₂ annual mean concentration by county, 1996.	19
Figure 2-19. Long-term ambient NO ₂ trend, 1977–1996.	20
Figure 2-20. Number of summer days, June–August with temperatures >90°, 1995 vs. 1996.	22
Figure 2-21. Trend in annual second daily maximum 1-hour O ₃ concentrations, 1987–1996.	23
Figure 2-22. O ₃ second daily maximum 1-hour concentration trends by location, 1987–1996.	23
Figure 2-23. Comparison of actual and meteorologically adjusted ozone trends, 1987–1996 (composite average of 99th percentile 1-hr daily max concentration).	24
Figure 2-24. Highest O ₃ second daily maximum concentration by county, 1996.	24
Figure 2-25. Long-term trend in second daily maximum 1-hour O ₃ concentrations, 1977–1996.	25
Figure 2-26. National total VOC emissions trend, 1987–1996.	25
Figure 2-27. VOC emissions by source category, 1996.	26
Figure 2-28. Trend in 2nd max 1-hr vs. 4th max 8-hr ozone concentrations, 1987–1996.	28
Figure 2-29. Trend in annual mean PM ₁₀ concentrations, 1988–1996.	30
Figure 2-30. National PM ₁₀ emissions trend, 1988–1996 (traditionally inventoried sources only).	31
Figure 2-31. PM ₁₀ annual mean concentration trends by location, 1988–1996.	31
Figure 2-32. PM ₁₀ emissions from traditionally inventoried source categories, 1996.	32
Figure 2-33. Total PM ₁₀ emissions by source category, 1996.	32
Figure 2-34. Highest second maximum 24-hour PM ₁₀ concentration by county, 1996.	33
Figure 2-35. PM ₁₀ trend in the average 99th percentile PM ₁₀ concentration, 1988–1996.	34
Figure 2-36. Highest second maximum 24-hour SO ₂ concentration by county, 1996.	38
Figure 2-37. Trend in annual mean SO ₂ concentrations, 1987–1996.	39

Figure 2-38. National total SO ₂ emissions trend, 1987–1996.	39
Figure 2-39. SO ₂ emissions by source category, 1996.	40
Figure 2-40. SO ₂ annual mean concentration trend by location, 1987–1996.	40
Figure 2-41. Long-term ambient SO ₂ trend, 1977–1996.	41
Figure 3-1. Range of best and worst conditions at Acadia, Great Smoky Mountains, and Grand Canyon national parks, 1992–1995.	44
Figure 3-2. Long-term trend for 75th percentile light extinction coefficient from airport visual data (July–September).	45
Figure 3-3. IMPROVE visibility monitoring network 30 sites with data for the period 1988–present.	46
Figure 3-4a. Total light extinction trends for eastern Class I areas.	47
Figure 3-4b. Total light extinction trends for western Class I areas.	47
Figure 3-5a. Light extinction due to sulfate in eastern Class I areas.	48
Figure 3-5b. Light extinction due to sulfate in western Class I areas.	48
Figure 3-6a. Light extinction due to organic carbon in eastern Class I areas.	48
Figure 3-6b. Light extinction due to organic carbon in western Class I areas.	48
Figure 3-7a. Average aerosol light extinction in eastern Class I areas.	49
Figure 3-7b. Average aerosol light extinction in western Class I areas.	49
Figure 3-8. Annual average light extinction (Mm ⁻¹), 1992–1995 IMPROVE data.	50
Figure 3-9. Annual average visibility impairment in deciviews relative to pristine conditions of deciviews = 0, 1992–1995 IMPROVE data.	51
Figure 3-10. Shenandoah National Park on clear and hazy days, and the effect of adding 10 µg/m ³ fine particles to each.	51
Figure 4-1. PAMS percent of total number of ozone nonattainment areas and 1996 ozone exceedance days (total number of original classified and section 185a ozone nonattainment areas = 118; total number of 1996 exceedance days in original nonattainment areas = 361.)	54
Figure 4-2. Comparison of actual and meteorologically adjusted ozone trends—PAMS metropolitan areas versus non-PAMS areas, 1987–1996 (composite average of 99th percentile 1-hr. daily max. conc.)	58
Figure 5-1. Total national HAP emissions by source type, 1993. (tons per year).	62
Figure 5-2. HAP emissions by state, 1993 (tons/year).	62
Figure 5-3. MACT source categories.	66
Figure 5-4. Emissions of 40 potential section 112(k) HAPs by source type (tons/year).	68
Figure 5-5. Emissions of 40 potential section 112(k) HAPs by urban and rural classification (tons/year).	69
Figure 6-1. Location of nonattainment areas for criteria pollutants.	73
Figure 6-2. Classified ozone nonattainment areas.	74
Figure 7-1. Number of days with PSI values > 100, as a percentage of 1987 value.	77
Figure A-1. (Multiple NA areas within a larger NA area) Two SO ₂ areas inside the Pittsburgh–Beaver Valley ozone NA. Counted as one NA area.	115
Figure A-2. (Overlapping NA areas) Searles Valley PM ₁₀ NA partially overlaps the San Joaquin Valley ozone NA. Counted as two NA areas.	115
Figure B-1. Carbon monoxide monitoring network, 1996.	150
Figure B-2. Lead monitoring network, 1996.	150
Figure B-3. Nitrogen dioxide monitoring network, 1996.	151
Figure B-4. Ozone monitoring network, 1996.	151
Figure B-5. PM ₁₀ monitoring network, 1996.	152
Figure B-6. Sulfur dioxide monitoring network, 1996.	152

Tables

Table 1-1.	Percent Change in National Air Quality Concentrations and Emissions, 1987–1996	1
Table 1-2.	Long-term Percent Change in National Air Quality Concentrations and Emissions	3
Table 2-1.	NAAQS in Effect in 1996	7
Table 3-1.	Summary of Class I Area Trend Analysis	50
Table 3-2.	IMPROVE Sites With Potential Upward Trends	50
Table 4-1.	Metropolitan Areas Requiring PAMS	53
Table 4-2.	PAMS Target List of VOCs	54
Table 4-3.	PAMS Targeted VOCs Ranked by Mean 6–9 am Concentration, Summer 1996	56
Table 4-4.	Number of Ozone NAAQS Exceedance Days, by PAMS Area	57
Table 4-5.	Summary of Changes in Summer Mean Concentrations for Ozone, NO _x , and Selected VOCs, 1995–1996 and 1994–1996	59
Table 5-1.	Top 20 Sources of 1993 Toxic Emissions of Hazardous Air Pollutants	63
Table 5-2.	Summary of Changes in Mean Concentration for HAPs Measured as a Part of the PAMS Program (24-hour measurements), 1994–1996*	64
Table 5-3.	Comparison of Loading Estimates for the Great Lakes.....	64
Table 5-4.	List of Potential 112(k) HAPs	68
Table 7-1.	Summary of MSA Trend Analysis, by Pollutant	76
Table 7-2.	Pollutant Standards Index Values with Pollutant Concentration, Health Descriptors, and PSI Colors	77
Table A-1.	National Air Quality Trends Statistics for Criteria Pollutants, 1987–1996	80
Table A-2.	National Carbon Monoxide Emissions Estimates, 1987–1996 (thousand short tons)	82
Table A-3.	National Lead Emissions Estimates, 1987–1996 (short tons)	83
Table A-4.	National Nitrogen Oxides Emissions Estimates, 1987–1996 (thousand short tons)	84
Table A-5.	National Volatile Organic Compounds Emissions Estimates, 1987–1996 (thousand short tons)	85
Table A-6.	National Particulate Matter (PM ₁₀) Emissions Estimates, 1987–1996 (thousand short tons)	86
Table A-7.	Miscellaneous and Natural PM ₁₀ Emissions Estimates, 1987–1996 (thousand short tons)	86
Table A-8.	National Sulfur Dioxide Emissions Estimates, 1987–1996 (thousand short tons)	87
Table A-9.	National Long-Term Air Quality Trends, 1977–1996	88
Table A-10.	National Air Quality Trends Statistics by Monitoring Location, 1987–1996	89
Table A-11.	Maximum Air Quality Concentrations by County, 1996	90
Table A-12.	Trends From IMPROVE Monitoring Sites, 1988–1995	104
Table A-13.	Condensed Nonattainment Areas List(a)	112
Table A-14.	Maximum Air Quality Concentrations by Metropolitan Statistical Area, 1996	116
Table A-15.	Metropolitan Statistical Area Air Quality Trends, 1987–1996	123
Table A-16.	Number of Days with PSI Values Greater Than 100 at Trend Sites, 1987–1996, and All Sites in 1996	144
Table A-17.	(Ozone only) Number of Days with PSI Values Greater Than 100 at Trend Sites, 1987–1996, and All Sites in 1996	146
Table A-18.	Total Number of Days with PSI Values Greater Than 100 at Trend Sites—Summary, 1987–1996	148
Table B-1.	Number of Ambient Monitors Reporting Data to AIRS	149

Acronyms

AIRS	Aerometric Information Retrieval System
CAA	Clean Air Act
CAAA	Clean Air Act Amendments
CARB	California Air Resources Board
CASAC	Clean Air Scientific Advisory Committee
CEMs	Continuous Emissions Monitors
CFR	Code of Federal Regulations
CO	Carbon Monoxide
CMSA	Consolidated Metropolitan Statistical Area
DST	Daylight Savings Time
EPA	Environmental Protection Agency
GDP	Gross Domestic Product
HAPs	Hazardous Air Pollutants
IMPROVE	Interagency Monitoring of PROtected Environments
MACT	Maximum Achievable Control Technology
MARAMA	Mid-Atlantic Regional Air Management Association
MSA	Metropolitan Statistical Area
NAAQS	National Ambient Air Quality Standards
NAMS	National Air Monitoring Stations
NARSTO	North American Research Strategy for Tropospheric Ozone
NESCAUM	Northeast States for Coordinated Air Use Management
NMOC	Non-Methane Organic Compound
NO ₂	Nitrogen Dioxide
NO _x	Nitrogen Oxides
NTI	National Toxics Inventory
O ₃	Ozone
OTAG	The Ozone Transport Assessment Group
PAHs	Polyaromatic Hydrocarbons
PAMS	Photochemical Assessment Monitoring Stations
Pb	Lead
PCBs	Polychlorinated Biphenyls
PM ₁₀	Particulate Matter of 10 micrometers in diameter or less
PM _{2.5}	Particulate Matter of 2.5 micrometers in diameter or less
POM	Polycyclic Organic Matter
ppm	Parts Per Million
PSI	Pollutant Standards Index
RFG	Reformulated Gasoline
SLAMS	State and Local Air Monitoring Stations
SNMOC	Speciated Non-Methane Organic Compound
SO ₂	Sulfur Dioxide
SO _x	Sulfur Oxides
TRI	Toxic Release Inventory
TSP	Total Suspended Particulate
VMT	Vehicle Miles Traveled
VOCs	Volatile Organic Compounds

